Spatial and temporal patterns of larval dispersal in a coral-reef fish metapopulation: evidence of variable reproductive success

TIMOTHY J. PUSACK,*† MARK R. CHRISTIE,*‡ DARREN W. JOHNSON,*§ CHRISTOPHER D. STALLINGS*† and MARK A. HIXON*¶

*Department of Integrative Biology, Oregon State University, Corvallis, OR 97331-2914, USA, †College of Marine Science, University of South Florida, St. Petersburg, FL 33701-5016, USA, ‡Department of Biological Sciences and Department of Forestry and Natural Resources, Purdue University, West Lafayette, IN 47907, USA, §Department of Biological Sciences, California State University Long Beach, 1250 Bellflower Blvd, Long Beach, CA 90840, USA, ¶Department of Biology, University of Hawai'i at Manoa, Honolulu, HI 96822-2216, USA

Abstract

Many marine organisms can be transported hundreds of kilometres during their pelagic larval stage, yet little is known about spatial and temporal patterns of larval dispersal. Although traditional population-genetic tools can be applied to infer movement of larvae on an evolutionary timescale, large effective population sizes and high rates of gene flow present serious challenges to documenting dispersal patterns over shorter, ecologically relevant, timescales. Here, we address these challenges by combining direct parentage analysis and indirect genetic analyses over a 4-year period to document spatial and temporal patterns of larval dispersal in a common coral-reef fish: the bicolour damselfish (Stegastes partitus). At four island locations surrounding Exuma Sound, Bahamas, including a long-established marine reserve, we collected 3278 individuals and genotyped them at 10 microsatellite loci. Using Bayesian parentage analysis, we identified eight parent-offspring pairs, thereby directly documenting dispersal distances ranging from 0 km (i.e., self-recruitment) to 129 km (i.e., larval connectivity). Despite documenting substantial dispersal and gene flow between islands, we observed more self-recruitment events than expected if the larvae were drawn from a common, well-mixed pool (i.e., a completely open population). Additionally, we detected both spatial and temporal variation in signatures of sweepstakes and Wahlund effects. The high variance in reproductive success (i.e., 'sweepstakes') we observed may be influenced by seasonal mesoscale gyres present in the Exuma Sound, which play a prominent role in shaping local oceanographic patterns. This study documents the complex nature of larval dispersal in a coral-reef fish, and highlights the importance of sampling multiple cohorts and coupling both direct and indirect genetic methods in order disentangle patterns of dispersal, gene flow and variable reproductive success.

Keywords: larval connectivity, parentage, self-recruitment, sweepstakes effects, Wahlund effect *Received 14 June 2013; revision received 9 May 2014; accepted 21 May 2014*

Introduction

Many marine animals have a biphasic life cycle that includes a pelagic larval stage followed by comparatively sedentary juvenile and adult stages. One

Correspondence: Timothy J. Pusack, Fax: (541) 737-0501; E-mail: pusackt@science.oregonstate.edu consequence of this ubiquitous life history strategy is that ocean currents can potentially transport pelagic larvae hundreds of kilometres from where they were spawned (Doherty *et al.* 1995; Weersing & Toonen 2009). Many such species therefore exist within marine metapopulations, where local populations of demersal juveniles and adults are connected by the exchange of planktonic larvae (Kritzer & Sale 2004). However, the pattern and magnitude of larval dispersal depends on a complex interplay among ocean currents (Gilg & Hilbish 2003; Cowen & Sponaugle 2009; White *et al.* 2010), larval behaviour (Leis & McCormick 2002; Gerlach *et al.* 2007), pelagic larval duration (Shanks 2009; Weersing & Toonen 2009; Faurby & Barber 2012) and physiological condition (Booth & Hixon 1999; Grorud-Colvert & Sponaugle 2011; Johnson *et al.* 2014). Variation in one or more of these factors can substantially affect both the spatial and temporal patterns of larval dispersal.

Stochastic processes associated with larval dispersal can lead to high larval mortality and may result in variable reproductive success where only a small proportion of adults produce the majority of larvae that survive through their pelagic larval phase (Hedgecock & Pudovkin 2011). Hedgecock (1994a,b) called this phenomenon 'sweepstakes reproductive success', because only a few adults would be 'big winners' in terms of reproductive success. Hallmark characteristics of sweepstakes reproductive success include genetically distinct cohorts and decreased genetic variation and genetic diversity when recruit populations are compared with adult populations (Hedgecock 1994a,b; Hedgecock & Pudovkin 2011). Variable reproductive success can also cause a Wahlund effect, characterized by an excess of homozygotes (Wahlund 1928; Hedrick 2005b). When this phenomenon occurs, genetically distinct cohorts accumulate within a single local population, creating an admixture of genetically differentiated individuals. Evidence of sweepstakes reproductive success and the Wahlund effect indicate spatial as well as temporal variation in larval sources. Documenting both patterns of larval dispersal and variable reproductive success will inform our understanding as to where marine populations lie along the continuum between demographically open (i.e., dependent upon larval connectivity) or closed (i.e., replenished by self-recruitment).

Researchers have directly documented larval dispersal by employing techniques such as otolith microchemistry (Swearer et al. 1999; Miller & Shanks 2004), chemical tagging of individuals (Jones et al. 1999, 2005), and increasingly, genetic parentage analysis (Jones et al. 2005; Planes et al. 2009; Christie et al. 2010a,b; Harrison et al. 2012; Saenz-Agudelo et al. 2012). Studies using these methods have documented both self-recruitment (Jones et al. 2005; Almany et al. 2007; Christie et al. 2010a) and larval connectivity (Planes et al. 2009; Christie et al. 2010b; Harrison et al. 2012). However, to date, no study has applied both direct (e.g. parentage analysis) and indirect (e.g. Wright's F statistics) genetic methods to monthly cohorts sampled over multiple years. Results from such broad-scale studies are critical for understanding marine metapopulation dynamics (Selkoe et al. 2006; Siegel et al. 2008; Watson et al. 2012),

© 2014 John Wiley & Sons Ltd

informing the design of marine reserve networks (Palumbi 2003; McCook *et al.* 2009), and enhancing fisheries management (Gell & Roberts 2003; Fogarty & Botsford 2007).

In this study, we coupled both direct parentage analysis and indirect genetic methods to characterize patterns of larval dispersal in a large marine metapopulation in Exuma Sound, Bahamas, over a 4-year period. Exuma Sound is a semi-enclosed basin characterized by a hydrodynamically complex environment that is punctuated with vigorous, semi-permanent and mesoscale gyres (Colin 1995; Stockhausen & Lipcius 2001). Depending upon their structure and dynamics, these gyres may facilitate connectivity between islands located on opposite sides of the Sound, may cause selfrecruitment within an island, or may simply create high mortality for larvae via entrainment and subsequent advection from suitable habitat. For example, simulated oceanographic models of spiny lobster (Panulirus argus) dispersal combined with abundance data found spatially variable levels of larval supply at locations throughout the Exuma Sound (Lipcius et al. 1997; Stockhausen et al. 2000; Stockhausen & Lipcius 2001). These results indicated that different hydrodynamic regimes of the gyres could affect the degree of connectivity between and self-recruitment within local populations in the Exuma Sound.

Our objective was to quantify temporal and spatial patterns of larval dispersal and variable reproductive success in bicolour damselfish (Stegastes partitus). We addressed four questions: (i) How consistent are the spatial and temporal patterns of connectivity and selfrecruitment in Exuma Sound? (ii) Given the low levels of genetic differentiation previously documented in bicolour damselfish (Purcell et al. 2009; Christie et al. 2010a), is there greater evidence for connectivity between sites than self-recruitment within sites? (iii) Are directly documented patterns of larval dispersal congruent with those inferred with indirect genetic methods? and (iv) Is there evidence for high variance in reproductive success? To answer these questions, we sampled adults and recruits (i.e., individuals that had recently settled from the plankton) of bicolour damselfish from multiple locations within a regional metapopulation and coupled parentage analyses with calculations of heterozygosity, genetic diversity and genetic differentiation to document patterns of larval dispersal and reproductive success.

Materials and methods

Study site and species

Approximately 500 km of continuous and shallow patch reefs encircle Exuma Sound, a semi-enclosed,

deep ocean basin (>1500 m deep) bordered by the Grand Bahamas Bank (<5 m deep) to the west and north, and the Atlantic Ocean to the east and south (Fig. 1). Circulation is partially self-contained, and consists of predominantly northwesterly along-shore currents (~20 cm/s; Colin 1995) that are punctuated by mesoscale gyres that last from 10 to 30 days and can reach a depth of 200 m (Lipcius *et al.* 1997; Stockhausen & Lipcius 2001).

The bicolour damselfish is a common inhabitant of coral reefs throughout the tropical Western Atlantic and greater Caribbean region (Humann & Deloach 2002). Juveniles and adults are highly site-attached and tend to stay within small (1-3 m²) home ranges (Myrberg 1972; Schmale 1981; Hixon et al. 2012). Bicolour damselfish do not form long-term mating pairs; instead, territorial males guard demersal nests and may receive eggs from multiple females. Females also mate with multiple males and will travel to male territories up to 10 m away when spawning (Knapp & Warner 1991; Cole & Sadovy 1995; Johnson et al. 2010). Spawning activity follows the lunar cycle and overall reproductive effort peaks during the summer months (Hixon et al. 2012). Eggs hatch after ~3.5 days of benthic development (Schmale 1981; Johnson et al. 2010). The pelagic larval duration (PLD) of bicolour damselfish ranges from 24 to 40 days (mean \approx

30 days, Wellington & Victor 1989) before late-stage larvae settle to suitable coral-reef habitat and recruit to the local population (Sponaugle & Cowen 1996). This PLD is typical of many tropical marine fishes (Victor and Wellington 2000, Lester and Ruttenberg 2005). In Exuma Sound, bicolour damselfish mature at 6-7 cm total length (TL) and live an average of 0.88 years, with a maximum recorded lifespan of 3.6 years (Hixon et al. 2012). Previous work on bicolour damselfish dispersal in Exuma Sound directly documented selfrecruitment through parentage analysis at two islands (Christie et al. 2010a). Christie et al. (2010a) also provided evidence for sweepstakes reproductive success, whereby the highly variable reproductive success of adults results in smaller genetic variation and diversity in each cohort (Hedgecock 1994a,b; Hedgecock & Pudovkin 2011).

Sample collection

From 2005 to 2008, we collected tissue samples from 3278 bicolour damselfish from three sites adjacent to each of four islands (12 sites total) encompassing Exuma Sound: Cat Island, Eleuthera, Exuma Cays Land and Sea Park and Lee Stocking Island (Fig. 1, the general locations are shown for the sample sites, which were all within 8 km of each other for each island). In



Fig. 1 (a) Adult bicolour damselfish, Stegastes partitus (courtesy of David Nasser). (b) Map of Exuma Sound, with sampled islands and corresponding adult (A) and recruit (R) sample sizes over all years. The arrows indicate the predominant currents in Exuma Sound, including seasonal mesoscale gyres (thick, grey swirls) and the prevailing northwesterly Antilles Current (thin, grey, dashed arrows). The yellow star represents a marine reserve study area and black circles are unprotected study areas. ECLSP, Exuma Cays Land and Sea Park; LSI, Lee Stocking Island. (c) Near-surface geostrophic currents in Exuma Sound computed from conductivity, temperature and depth (CTD) observations (Stockhausen & Lipcius 2001). Notice that the gyres are closer to ECLSP, Eleuthera and Cat Island than to LSI. (d) Satellite image of one of the northern seasonal gyres adjacent to Eleuthera (courtesy of Larry Ammann, University of Texas).

2005, samples were collected during a single summer visit at each of the four islands (Christie *et al.* 2010a). From 2006 to 2008, we collected adult and recruit samples at the same sites in June, July and August to correspond with the peak summer spawning and recruitment events. Because of the short interval between visits, and by collecting only fish \leq 30 mm (TL), it was reasonable to assume that all of the sampled recruits in each month belonged to the same cohort (Sponaugle & Cowen 1996).

To facilitate collection with hand nets, SCUBA divers anaesthetized individual damselfish with a solution of 10% quinaldine and 90% methanol mixed with equal amounts of seawater. The anaesthetic solution was targeted at specific fish using squeeze bottles

to minimize exposure to other marine life. We clipped a small portion of the pelvic fins from adults (>5 cm TL, n = 1348) before releasing them at the collection site. Recently settled recruits (<3 cm TL, n = 1930) were euthanized and their caudal fins were preserved for subsequent analysis. All tissue samples were stored in a urea-based solution (10 mM Tris, 125 mM NaCl, 10 mM EDTA, 1% SDS, 8 M urea, pH adjusted to 7.5 with HCL). The numbers of adults sampled varied by island and year based on the number of new adults that became established in each of the sample sites. We haphazardly collected tissue samples from about 50 recruits at each monthly visit to each island. Thus, we sampled approximately 150 recruits from each island per summer (Table 1).

 Table 1 Summary statistics of local populations of bicolour damselfish

Island	Year	Ν	$N_{\rm a}$	k	$F_{\rm IS}$	H _o	$H_{\rm e}$	HWE	% LD
Adults									
Cat	2005	46	23	19	0.102	0.834	0.928	3/7	0
Cat	2006	115	26	18	0.103	0.803	0.895	3/10	0
Cat	2007	123	24	18	0.088	0.814	0.893	3/10	0
Cat	2008	147	26	18	0.076	0.823	0.890	3/10	0
ECLSP	2005	47	22	19	0.060	0.874	0.928	2/7	0
ECLSP	2006	63	20	17	0.104	0.798	0.891	2/10	0
ECLSP	2007	59	21	18	0.097	0.811	0.897	2/10	0
ECLSP	2008	64	22	18	0.011	0.789	0.889	2/10	0
Eleuthera	2005	49	22	19	0.052	0.887	0.936	0/7	0
Eleuthera	2006	43	20	18	0.930	0.809	0.891	2/10	0
Eleuthera	2007	107	24	18	0.091	0.810	0.890	2/10	0
Eleuthera	2008	105	25	18	0.115	0.790	0.892	2/10	0
LSI	2005	43	20	18	0.036	0.899	0.932	0/7	0
LSI	2006	104	28	19	0.120	0.856	0.932	3/7	0
LSI	2007	108	19	18	0.091	0.817	0.893	2/10	0
LSI	2008	125	26	18	0.104	0.799	0.892	3/10	0
Recruits									
Cat	2005	44	22	18	0.090	0.843	0.925	2/7	0
Cat	2006	152	29	18	0.104	0.801	0.894	5/10	0
Cat	2007	155	26	18	0.104	0.799	0.892	6/10	0
Cat	2008	147	26	18	0.110	0.794	0.893	3/10	0
ECLSP	2005	45	22	18	0.014	0.916	0.929	1/7	0
ECLSP	2006	108	25	18	0.091	0.812	0.894	3/10	0
ECLSP	2007	115	25	18	0.094	0.810	0.894	2/10	0
ECLSP	2008	129	24	18	0.107	0.796	0.890	2/10	0
Eleuthera	2005	37	21	19	0.049	0.883	0.927	1/7	4.4
Eleuthera	2006	163	26	18	0.077	0.823	0.892	2/10	0
Eleuthera	2007	145	26	18	0.110	0.791	0.888	4/10	0
Eleuthera	2008	142	26	18	0.121	0.784	0.891	5/10	0
LSI	2005	74	25	19	0.081	0.858	0.933	3/7	0
LSI	2006	162	27	18	0.104	0.804	0.897	3/10	0
LSI	2007	162	28	18	0.124	0.781	0.891	5/10	0
LSI	2008	150	25	18	0.147	0.758	0.888	5/10	0

ECLSP, Exuma Cays Land and Sea Park; LSI, Lee Stocking Island. Column headings are as follows: N, sample size, $N_{a'}$ mean number of alleles per locus, k, mean allelic richness (corrected for smallest sample size), F_{IS} , Wright's inbreeding coefficient, $H_{o'}$ observed heterozygosity, $H_{e'}$ expected heterozygosity, HWE, the number of loci that deviate from Hardy–Weinberg Equilibrium/total number of loci, % LD, percentage of loci pairs in linkage disequilibrium.

DNA extraction and microsatellite typing

DNA was extracted using a protocol optimized for the urea-based storage solution (Sambrook & Russell 2001; see Christie et al. 2010a for details). Samples from 2005 (n = 385) were previously extracted and genotyped at seven microsatellite loci (Christie et al. 2010a), and because adults sampled in 2006 from Lee Stocking Island (n = 104) were needed for quantitative genetic studies (Johnson et al. 2010, 2011), they were also genotyped at the same seven microsatellite loci: SpAAC33, SpAAC 41, SpAAC42, SpAAC47, SpAAT40, SpGATA16 and SpGAT-A40 (Williams et al. 2003). The remaining 2839 samples were genotyped at the above seven loci and three additional microsatellite loci: SpTG10, SpTG16 and SpGGA7 (Thiessen & Heath 2007). PCRs were optimized and multiplexed with the Qiagen Multiplex PCR Kit and contained 5.0 µL of Qiagen Multiplex mix, 1.0 µL of each primer and 2.0 µL of approximately 100 ng/µL DNA template for a total volume of 10 µL. The thermocycle profile consisted of an initial denaturing step at 95 °C for 15 min, followed by 30 cycles of 30 s at 94 °C, 90 s at 58 °C and 60 s at 72 °C. PCR products were screened on an ABI 3100 automated sequencer (Applied Biosystems) at the Center for Genome Research and Biocomputing at Oregon State University. We used GENOTYPER 3.7 to determine allele sizes. We reprocessed 19.5% (n = 640) of the individuals (from extraction through scoring) to check for errors in genotyping and to check individuals that were difficult to score at the most polymorphic loci (the error rate was 3.7%).

Parentage analyses

To directly document dispersal on an ecological timescale, we used a Bayesian parentage analysis (Christie 2010; Christie et al. 2013) to identify parent-offspring pairs. Briefly, this method calculates the posterior probability of any given pairwise comparison sharing alleles by chance, given the frequencies of the shared alleles. We compared the genotypes of all biologically plausible adults and all recruits to detect putative parent-offspring pairs. Using the extensive demographic data, we excluded adults that could not be possible parents (due to mortality in previous years) for each cohort of recruits. Pairs unlikely to share alleles by chance (i.e., posterior probability <0.05) after fully accounting for all possible pairwise comparisons were considered to be parentoffspring pairs. All putative pairs were re-analysed from DNA extraction through scoring at all loci to eliminate the possibility of laboratory errors. All pairs shared at least one allele at all loci. None of the identified parentoffspring pairs had missing data.

The possibility of pairs actually being some other first-order relative, that is full-siblings, was very

unlikely due to the size differences between adults and the offspring in each pair. For example, based on average growth rates in our study area (Hixon et al. 2012), it would take 430-750 days for a fish to grow to the sizes of the parents. Assuming that the instantaneous mortality rate of adults is 0.0045 per day (Hixon et al. 2012), the probability of an adult living long enough to have been the parents of both members of our putative parent-offspring pairs was low (0.034-0.144). We also used a simulation to evaluate the probability that full-siblings could have shared at least one allele at all loci, and thus exhibited the same allele matching pattern as parentoffspring pairs. We used the allele frequencies from the entire population and created genotypes in accordance with Hardy-Weinberg Equilibrium (HWE). Next, we randomly paired individuals and simulated full-sibling offspring based on expected patterns of Mendelian inheritance. We then counted how many full-siblings shared at least one allele at all loci (as did our putative pairs). Because we had eight parent-offspring pairs, we created eight full-sibling pairs (each pair with different parents) and then counted the number of pairs that shared an allele at each locus. This process was repeated 1000 times, and we calculated the probability of observing one full-sibling pair, two full-sibling pairs, three full-sibling pairs, etc. that shared at least one allele across all loci in our data set. We used the R statistical environment for these simulations (R Core Team 2012). Based on the number of shared alleles, the probability of one identified 'parent-offspring pair' being full-siblings was not trivial (P = 0.76). The probability of two or more pairs being full-siblings was low (P = 0.04), and the probability of three or more pairs being full-siblings was very unlikely (P = 0.001). These analyses suggest that, at most, one of the identified 'parent-offspring pairs' could have been full-siblings. However, this event was unlikely when considered within the context of the above life history information. Given the low probability that one fish lived long enough to produce a 6-cm adult and a 2.5 cm recruit, and the low probability of fullsiblings sharing one allele at each loci, it is unlikely that any of the documented parent-offspring pairs were fullsiblings or higher-order relatives.

Dispersal distances between identified parent and offspring were determined using both Euclidean distance and along-shore distance (the nearest along-shore path between the adult sample site and the recruit sample site following the fore-reef contour at a resolution of 1 km). Because bicolour damselfish do not move more than a few metres after settling to reef habitat (Hixon *et al.* 2012), the Euclidian distances between parent and offspring represent the minimum dispersal distances of planktonic larvae (i.e., the actual distances travelled by dispersing larvae may have been much greater if they did not follow a direct path).

Given the low genetic differentiation previously documented in this system (Christie et al. 2010a), it is possible that larvae from multiple source populations are well mixed while in the plankton. Because parent-offspring pairs record the starting and ending points for larval dispersal, we could test whether or not our observed pattern of larval dispersal (five self-recruitment events and three connectivity events) was consistent with homogenous mixing of larvae from each of the sampled populations. As a null hypothesis, we expected islands to contribute to a randomly mixed pool of larvae that supplies all islands with recruits. However, the probability of observing selfrecruitment or connectivity among the sites depends on the number of parents and offspring sampled at each site, and these values were unequal in our study. Using the Rstatistical environment (R Core Team 2012), we simulated a randomly mating population by creating simulated adults, represented by the numbers of adults we sampled at each island, and placed them together in a combined population. We next sampled this mixed population by the total number of parents we identified, eight individuals, and noted from which population they were sampled. Then, using the same procedure, we sampled simulated recruits and generated locations of parents and offspring expected in an open population, where larvae from each local population joined a well-mixed pool of larvae that contributed offspring to all local populations. We repeated this procedure 10 000 times and calculated the *P*-value as the proportion of simulations that identified equal or greater numbers of self-recruitment events to the one we observed for our empirical data. Using this framework, we tested whether the number of parent-offspring pairs sampled at the same sties (i.e., self-recruitment) was less than or greater than expected.

Population-genetic analyses

GENEPOP v. 4.0.11 (Raymond & Rousset 1995; Rousset 2008) was first employed to determine whether any locus pairs were in linkage disequilibrium, and to test for departures from HWE for adult and recruit samples collected each year at each island. For each measure, a total of 10 000 batches and 5000 iterations per batch were used to reduce the standard errors below 0.01. We tested for large allele dropout, null alleles and stuttering that could explain deviations from HWE using MICRO-CHECKER v2.2.3 (Van Oosterhout et al. 2004). We used ARLEQUIN v3.11 (Excoffier et al. 2005) to calculate observed and expected heterozygosities and FSTAT v2.9.3.2 (Goudet 2001) to calculate F_{IS} values, mean number of alleles per locus, mean allelic richness (k) and Nei's gene diversity (1987). Allelic richness was corrected for sample size of adult and recruit population at each island among years. To test for population

differentiation, we used a variety of metrics based on allele frequencies in each population. We first used FSTAT to calculate pairwise, unbiased F_{ST} , (Weir & Cockerham 1984) and we adjusted the corresponding P-values for multiple comparisons with a sequential Bonferroni procedure ($\alpha = 0.05$). We also used randomization tests within FSTAT (5000 permutations) to compare F_{ST} values between recruits pooled by year. For comparisons between groups with ten and seven loci, only the matching seven loci were used. Because highly polymorphic loci can artificially reduce F_{ST} values (Hedrick 2005a; Meirmans & Hedrick 2011), we also used SMOGD v1.2.5 (Crawford 2010) to calculate pairwise Dest values (Jost 2008) and GENODIVE (Meirmans & Van Tiernderen 2004) to calculate standardized versions similar to F_{ST} including multilocus G'_{ST} (Hedrick 2005a) and pairwise population G''_{ST} (Meirmans & Hedrick 2011).

Variable reproductive success

To test for the Wahlund effect, we compared the number of loci out of HWE due to a homozygous excess (averaged across cohorts) to cohort groups that were pooled together, both temporally (each island across all years) and spatially (each month across all islands). To examine the evidence for a temporal Wahlund effect, we compared the average number of loci that deviated from HWE due to homozygote excess for each recruit cohort at each island (averaged across months) to the samples pooled across all months and years at that island. To examine the evidence for a spatial Wahlund effect, we compared the average number of loci that deviated from HWE due to homozygote excess for each monthly recruit cohort (averaged across islands) to the samples pooled across all islands in a given month. If a Wahlund effect occurred, then the number of loci with a homozygous excess would be less in the cohort average compared with the pooled group. To test for sweepstakes reproduction, we first tested for genetic differences among each cohort using a randomization test (5000 permutations) in FSTAT to compare F_{ST} among pooled recruit samples by islands and years. Second, we used randomization tests (5000 permutations) in FSTAT to test for differences in genetic variation (observed heterozygosity), genetic diversity (expected heterozygosity and allelic richness), and F_{IS} between recruits and adults sampled in each year.

Results

Parentage analysis

We detected eight parent-offspring pairs, two each year from 2005 to 2008. We documented five pairs that exhibited self-recruitment within an island (two of which were previously reported by Christie *et al.* 2010a) and three pairs that documented connectivity between islands (Fig. 2, Table 2). Remarkably, in three of the five self-recruitment events, the recruit fish returned to the exact reef from which it originated (Table 2). For the other two self-recruitment events, the recruits returned to a nearby site at the same island (<5 km away). The longest dispersal distance detected was 111.2 km in Euclidean distance, or 129.4 km in along-shore distance (Table 2). Given the sample sizes we collected and the number of parent-offspring pairs we detected, it is unlikely that we would have documented so many self-recruitment events, if the larvae were drawn from a common, well-mixed pool (randomization test, *P*-value = 0.03).

Population-level analyses

Across all adult and recruit populations grouped by yearly collection at each island, the mean number of alleles per population over all 10 loci ranged from 19 to 29. The mean allelic richness per population over all loci, calculated from a minimum sample size of 43 for adults and 37 for recruits, ranged from 17 to 19 alleles. Observed heterozygosity ranged from 0.758 to 0.916 (Table 1). Based on the MICRO-CHECKER analysis for deviations from HWE, homozygote excess was the most frequent cause of deviation. There was no evidence for

large allele drop-out, and we detected low average numbers of null alleles in our 10 loci: 2.4 [0.22 standard error or the mean (SEM)] in adult populations and 2.8 (0.18 SEM) in recruit populations. We suspect that a large proportion of homozygous genotypes are from population admixture (i.e., a Wahlund effect: see 'variable reproductive success' below).

Global F_{ST} was low and not significantly different from zero (95% confidence interval: 0.000-0.001). Pairwise comparisons of F_{ST} revealed low genetic differentiation across all years and islands for both adult and recruit populations, ranging from -0.0032 to 0.0046 (Table S1, Supporting information). The low F_{ST} values were not simply an artefact of the highly polymorphic loci. Similar metrics that account for polymorphic loci were also relatively low: G'ST ranged from -0.003 to 0.005, $G''_{\rm ST}$ ranged from -0.045 to 0.065, and $D_{\rm est}$ ranged from -0.016 to 0.057 with a mean for each estimator of 0.0004, 0.006 and 0.004, respectively. Nevertheless, after correcting for multiple comparisons, there were 21 significant F_{ST} pairwise comparisons among adult populations out of the 120 total possible comparisons among islands across years. Lee Stocking Island was one of the populations involved in the majority of those significant comparisons (86%; Table S1, Supporting information). Comparisons of adult to recruit populations across all years and islands revealed 20.3% of the pairwise comparisons of $F_{\rm ST}$ values to be



Fig. 2 Parent-offspring pairs of bicolour damselfish documented in each of 4 years. Orange solid arrows indicate larval connectivity and white dashed arrows show self-recruitment. The yellow star is the marine reserve study area, and black circles are unprotected study areas. ECLSP, Exuma Cays Land and Sea Park; LSI, Lee Stocking Island.

				Distance (km)		
Adult island (spawning island)	Offspring island (recruitment island)	Year	Month	Euclidean	Along shore	P-value
Eleuthera	Eleuthera	2005	July	0	0	0.0360
ECLSP	ECLSP	2005	July	0	0	0.0110
Cat	Cat	2006	August	0	0	0.0116
ECLSP	Eleuthera	2006	August	61.1	129.4	0.0433
Cat	Cat	2007	June	3.1	3.2	0.0259
Eleuthera	Cat	2007	August	111.2	129.0	0.0377
Cat	Eleuthera	2008	July	111.2	129.0	0.0332
ECLSP	ECLSP	2008	July	4.4	4.7	0.0136

Table 2 Euclidean and along-shore distance between members of each parent-offspring pair of bicolour damselfish

ECLSP, Exuma Cays Land and Sea Park.

Along-shore distance was measured by tracing the contour of the 20-m isobaths with a resolution of 1 km. The '*P*-value' is the posterior probability of the pair being false given the shared allele frequencies (Christie 2010). Notice that three offspring returned to the exact same reef inhabited by their parents (distance = 0).

significant (Table S2, Supporting information). Pairwise comparisons among all recruits among islands across years revealed that 22 of 120 pairwise comparisons were significant (18.3%) at an alpha equal to 0.05 after a sequential Bonferroni correction, and populations sampled at Lee Stocking Island were one of the pairs involved in 68% of the significant comparisons (Table S3, Supporting information). Finally, genetic differentiation among cohorts of recruits was significant when the following years were compared (FSTAT randomization test): 2005 vs. 2006 (*P*-value = 0.0056); 2005 vs. 2008 (*P*-value = 0.0146); 2006 vs. 2007 (*P*-value = 0.0034); and 2007 vs. 2008 (*P*-value = 0.0252; 2005 vs. 2007 was marginally significant; *P*-value = 0.105).

Variable reproductive success

We found evidence for both temporal and spatial Wahlund effects. We compared the average number of loci that deviated from HWE for each recruit cohort at each island (averaged across months) to recruit samples pooled across all months and years at that island. We found that the pooled estimate had considerably more loci out of HWE due to homozygote excess (Fig. 3a), indicating a temporal Wahlund effect (i.e., samples across time were more different from one another than samples within the same month). We also compared the average number of loci that deviated from HWE for each monthly recruit cohort (averaged across islands) to the samples pooled across all islands in a given month and found that the pooled estimates had considerably more loci out of HWE due to homozygote excess (Fig. 3b), indicating a spatial Wahlund effect (i.e., samples across islands in a given month were more different from one another than samples within a site).

As predicted by the sweepstakes hypothesis, we detected evidence for lower genetic variance and diversity in recruit populations compared with adult populations. The randomization tests in FSTAT revealed lower observed heterozygosity (P = 0.016) and expected heterozygosity (P = 0.019) in recruits compared with adults (Fig. 4a, b). Also the numbers of alleles (standardized by sample size) present in recruit populations were lower than adult populations (P = 0.032; Fig. 4c). Finally, we also documented a positive and significantly higher $F_{\rm IS}$ in recruit populations compared with adult populations (P = 0.034; Fig. 4d; again using FSTAT).

Discussion

We found direct evidence for both connectivity and self-recruitment in a marine metapopulation of bicolour damselfish. Expanding the temporal scale of Christie et al. (2010a) by 3 years, we documented dispersal events ranging up to ~130 km, which were previously inferred indirectly, but not directly quantified. Importantly, we also found that a long-established marine reserve, the Exuma Cays Land and Sea Park, not only was self-seeding (i.e., self-recruitment was documented twice at this island), but also generated larvae that seeded unprotected areas. While Lipcius et al. (2001) had inferred indirectly that this marine reserve is capable of both self-recruitment and exporting larvae to other islands, we provide the first direct evidence. Additionally, we documented a spatial Wahlund effect, which suggests that different sets of adults contributed to the offspring sampled at each island within a given month. We also documented to a lesser extent a temporal Wahlund effect, which suggests that each monthly cohort within a site originated from different sets of



Fig. 3 Temporal and spatial Wahlund effects. Each solid symbol is the number of loci that deviate from Hardy–Weinberg Equilibrium (HWE) averaged across cohorts [±1 standard error of the mean (SEM)]. Each open symbol is the number of loci that deviate from HWE when all cohorts are pooled. (a) Temporal Wahlund effect where closed symbols represent recruit populations averaged across months at each island and open symbols represent all recruit cohorts pooled at a given island. (b) Spatial Wahlund effect where closed symbols represent the number of loci out of HWE averaged across islands for recruits in a given month and open symbols represent recruits pooled across islands in the same month.

parents. Moreover, because we observed lower heterozygosity (observed and expected) and allelic richness in recruit cohorts compared with adult populations, it appears that sweepstakes reproduction was also prevalent in this system.

While many studies have indirectly inferred connectivity and self-recruitment, few studies have addressed the question of temporal variability of these events using direct methods (but see Saenz-Agudelo *et al.* 2012). To our knowledge, our study is the first to directly document larval dispersal patterns over 4 years, using multiple cohorts within each year. Our findings suggest that studies based on a single year may not sufficiently capture the variation in patterns of larval dispersal that can exist in marine systems. Conducting multi-year studies that sample multiple cohorts within each year to describe the variability in dispersal will facilitate effective planning for fisheries management and enhance the design and placement of marine reserves.

The pattern of gene flow inferred from F_{ST} suggested similar patterns of dispersal compared with the parentage analysis: the greatest number of significantly different pairwise comparisons of population structure was found at Lee Stocking Island, where no parent-offspring pairs were documented. The fact that one island, Lee Stocking Island, had the majority of genetic difference compared with itself as well as to other locations, suggests that dispersal may not be equal throughout the Exuma Sound. Similar levels of genetic differentiation have been documented in bicolour damselfish populations from other regions (e.g., Lacson 1992; Ospina-Guerrero *et al.* 2008; Hepburn *et al.* 2009; Hogan *et al.* 2010; Salas *et al.* 2010) and across the entire Caribbean region (Purcell *et al.* 2009). The temporal stability of genetic differences involving Lee Stocking Island means that these differences were not simply due to sampling error (Waples 1998), and may indicate that the patterns observed on an ecological timescale correspond to patterns on an evolutionary timescale.

Similar to other studies in marine systems, we detected a Wahlund effect despite low FST values (Tracey et al. 1975; Johnson & Black 1984; Addison & Hart 2004; Levitan 2005). These studies attributed localized genetic differentiation to be the mechanism causing the observed pattern. In our system, it is likely that sweepstakes reproduction was the mechanism resulting in the Wahlund effect. Our findings concur with previous work in this system by Christie et al. (2010a) that provided evidence of sweepstakes reproductive success. Because evidence of the sweepstakes reproduction in bicolour damselfish was also found from studies in the Mesoamerican Barrier Reef System (Hepburn et al. 2009; Villegas-Sanchez et al. 2010), this effect may be prevalent throughout the geographic range of this species. More generally, sweepstakes reproduction, and not necessarily restricted larval dispersal, may be responsible for patterns of genetic differentiation observed in many marine species.

The variable pattern of larval dispersal and reproductive success documented in this study has wide-ranging ecological and evolutionary implications. First, while variance in reproductive success is appreciated in many



terrestrial systems (Clutton-Brock 1988), the contribution of this variance is not well understood in marine systems. Second, many marine populations are characterized by high rates of dispersal (Levin 2006; Gaines et al. 2007) and gene flow (Waples 1998; Hedgecock et al. 2007). However, if there is high variance in reproductive success, then the resulting changes to allele frequency distributions (Hedgecock & Pudovkin 2011) can cause samples from populations to be genetically differentiated from one another even though there may be little to no correlation between dispersal and the documented genetic differentiation. Because this high variance in reproductive success is simply one form of genetic drift, and is thus random, there is often no relationship between geographic distance and genetic differentiation (sometimes termed 'chaotic genetic patchiness'; Johnson and Black 1982, Johnson & Black 1984). We suggest that there is nothing 'chaotic' about these observed patterns of genetic differentiation, and that by greatly extending the duration of sampling presented in Christie et al. (2010a), we have identified that (i) there is substantial dispersal in this system, as characterized by parents and offspring being as far as 129 km apart; and (ii) the signature of high gene flow in this system is continually eroded by high variance in reproductive success (i.e., sweepstakes reproduction). Thus, disentangling dispersal, gene flow and genetic drift can be achieved only by sampling multiple cohorts of recruits over multiple years or spawning cycles.

Although mechanisms for variance in reproductive success have rarely been documented, region-wide genetic homogeneity may be eroded by the predominant mesoscale gyres that commonly occur in Exuma Sound (Colin 1995; Stockhausen & Lipcius 2001). Depending upon their characteristics, each gyre may promote connectivity or self-recruitment and may also entrain larvae and advect them away from suitable habitat, causing high levels of larval mortality. Using biophysical models (reviewed by Werner et al. 2007) in our system could help to explain the mechanisms underlying the observed patterns. For example, at islands where offshore gyres have been previously observed-Exuma Cays Land and Sea Park, Eleuthera, and Cat Island-we detected self-recruitment, whereas no parent-offspring pairs were observed at Lee Stocking Island, where gyres are less prevalent (Stockhausen & Lipcius 2001). However, the hypothesis that gyres facilitate self-recruitment remains to be tested. Moreover, seasonal and annual fluctuations in location, duration and strength of the gyres may further affect patterns of successful larval dispersal. Such variation in the gyres may dictate whether oceanic conditions are favourable for successful settlement of larvae (Hedgecock & Pudovkin 2011), and thus may be the primary mechanism driving sweepstakes reproduction in this system. The other hydrodynamic feature of Exuma Sound, the prevailing northwesterly Antilles Current, could have transported larvae from Cat Island to Eleuthera, and when this current reverses direction, which can happen for short periods of time (Colin 1995), the opposite pattern may have occurred. This possibility was corroborated by the fact that we detected larval connectivity in both directions between these islands.

This study illustrates the temporally and spatially complex nature of larval dispersal in a coral-reef fish. Here, we show that by sampling multiple cohorts and coupling both direct and indirect genetic methods, it is possible to disentangle patterns of dispersal, gene flow and variable reproductive success. The next steps towards a comprehensive understanding of marine metapopulation dynamics include quantifying larval retention and linking genetic analyses with local demographic, oceanographic and environmental data. Synthesis of these different fields will greatly advance our understanding of metapopulation dynamics and enhance fisheries management and conservation

Fig. 4 Comparisons of the average observed heterozygosity, gene diversities (expected heterozygosity and allelic richness—k) and F_{IS} between adult (triangles) and recruit (circles) bicolour damselfish.

efforts by identifying the spatial boundaries of marine populations.

Acknowledgements

We thank S. Arnold and M. Blouin for laboratory space and valuable statistical advice. J. Moye and M. Albins deserve special thanks for their immense help in the field, and R. Lamb, M. Cooke, M. Errand, and M. Head for assistance in the laboratory and/or the field. We are grateful to the staff of the Perry Institute for Marine Science, especially Brenda Gadd and Ikeman Black. Special thanks to J.F.H. Purcell for detailed advice regarding storage, extraction and amplification of damselfish samples. We are grateful to L. Harrington for graphic design assistance. This work was supported by grants to M.A. Hixon from the National Science Foundation (OCE-00-93976 and OCE-05-50709).

References

- Addison JA, Hart MW (2004) Analysis of population genetic structure of the green sea urchin (*Strongylocentrotus droebachiensis*) using microsatellites. *Marine Biol*ogy, **144**, 243–251.
- Almany GR, Berumen ML, Thorrold SR, Planes S, Jones GP (2007) Local replenishment of coral reef fish population in a marine reserve. *Science*, **316**, 742–744.
- Booth DJ, Hixon MA (1999) Food ration and condition affect early survival of the coral reef damselfish, *Stegastes partitus*. *Oecologia*, **121**, 364–368.
- Christie MR (2010) Parentage in natural populations: novel methods to detect parent-offspring pairs in large data sets. *Molecular Ecology Resources*, **10**, 115–128.
- Christie MR, Johnson DW, Stallings CD, Hixon MA (2010a) Self-recruitment and sweepstakes reproduction amid extensive gene flow in a coral-reef fish. *Molecular Ecology*, **19**, 1042–1057.
- Christie MR, Tissot BN, Albins MA *et al.* (2010b) Larval connectivity in an effective network of marine protected areas. *PLoS ONE*, **5**, 1–8.
- Christie MR, Tennessen JA, Blouin M (2013) Bayesian parentage analysis with systematic accountability of genotyping error, missing data and false matching. *Bioinformatics*, 29, 725–732.
- Clutton-Brock TH (1988) Reproductive Success: Studies of Individual Variation in Contrasting Breed Systems (ed. Clutton-Brock TH). University of Chicago Press, Chicago, Illinois.
- Cole KS, Sadovy Y (1995) Evaluating the use of spawning success to estimate reproductive success in a Caribbean reef fish. *Journal of Fish Biology*, **47**, 181–191.
- Colin PL (1995) Surface currents in Exuma Sound, Bahamas and adjacent areas with reference to potential larval transport. *Bulletin of Marine Science*, **56**, 48–57.
- Cowen RK, Sponaugle S (2009) Larval dispersal and marine population connectivity. *Annual Review of Marine Science*, 1, 443–466.
- Crawford NG (2010) SMOGD: software for the measurement of genetic diversity. *Molecular Ecology Resources*, **10**, 556–557.
- Doherty PJ, Planes S, Mather P (1995) Gene flow and larval duration in seven species of fish from the Great Barrier Reef. *Ecology*, **76**, 2373–2391.

- Excoffier L, Laval G, Schneider S (2005) ARLEQUIN ver. 3.0: an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online*, **1**, 47–50.
- Faurby S, Barber PH (2012) Theoretical limits to the correlation between pelagic larval duration and population genetic structure. *Molecular Ecology*, **21**, 3419–3432.
- Fogarty MJ, Botsford LW (2007) Population connectivity and spatial management of marine fisheries. *Oceanography*, **20**, 112–123.
- Gaines SD, Gaylord B, Gerber LR, Hastings A, Kinlan BP (2007) Connecting places the ecological consequences of dispersal in the sea. *Oceanography*, 20, 90–99.
- Gell FR, Roberts CM (2003) Benefits beyond boundaries: the fishery effects of marine reserves. *Trends in Ecology and Evolution*, **18**, 448–455.
- Gerlach G, Atema J, Kingsford MJ, Black KP, Miller-Sims V (2007) Smelling home can prevent dispersal of reef fish larvae. *Proceedings of the National Academy of Science*, **104**, 858– 863.
- Gilg MR, Hilbish TJ (2003) The geography of marine larval dispersal: coupling genetics with fin-scale physical oceanography. *Ecology*, 84, 2989–2998.
- Goudet J (2001) FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3). Available from http:// www.unil.ch/izea/softwares/fstat.html.
- Grorud-Colvert K, Sponaugle S (2011) Variability in water temperature affects trait-mediated survival of newly settled coral reef fish. *Oecologia*, **165**, 675–686.
- Harrison HB, Williamson DH, Evans RD *et al.* (2012) Larval export from marine reserves and the recruitment benefit for fish and fisheries. *Current Biology*, **22**, 1023–1028.
- Hedgecock D (1994a) Does variance in reproductive success limit effective population size of marine organisms? In: *Genetics and Evolution of Aquatic Organism* (ed. Beaumont AR), pp. 122–135. Chapman and Hall, London.
- Hedgecock D (1994b) Temporal and spatial genetic structure of marine animal populations in the California current. *Califor*nia Cooperative Oceanic Fisheries Investigations, 35, 73–78.
- Hedgecock D, Pudovkin AI (2011) Sweepstakes reproductive success in highly fecund marine fish and shellfish: a review and commentary. *Bulletin of Marine Science*, **8**, 971–1002.
- Hedgecock D, Barber PH, Edmands S (2007) Genetic approaches to measuring connectivity. *Oceanography*, 20, 70–79.
- Hedrick PW (2005a) A standardized genetic differentiation measure. Evolution, 59, 1633–1638.
- Hedrick PW (2005b) *Genetics of Populations*. Jones and Bartlett Publishers, Boston, Massachusetts.
- Hepburn RI, Sale PF, Dixon B, Heath DD (2009) Genetic structure of juvenile cohorts of bicolor damselfish (*Stegastes partitus*) along the Mesoamerican barrier reef: chaos through time. *Coral Reefs*, 28, 277–288.
- Hixon MA, Anderson TW, Buch KL *et al.* (2012) Density dependence and population regulation in marine fish: a large scale, long-term field manipulation. *Ecological Monographs*, 82, 467–489.
- Hogan JD, Thiessen RJ, Heath DD (2010) Variability in connectivity indicated by chaotic genetic patchiness within and among populations of a marine fish. *Marine Ecology Progress* Series, 417, 263–275.

- Humann P, Deloach N (2002) Reef Fish Identification: Florida, Caribbean, Bahamas (ed. Deloach N), 3rd edn. New World Publications, Jacksonville, Florida.
- Johnson MS, Black R (1984) The Wahlund effect and the geographical scale of variation in the intertidal limpet *Siphonaria* sp. *Marine Biology*, **79**, 295–302.
- Johnson MS, Black R (1982) Chaotic genetic patchiness in an intertidal limpet, *Siphonaria* sp. *Marine Biology*, **70**, 157–164.
- Johnson DW, Christie MR, Moye J (2010) Quantifying evolutionary potential of marine fish larvae: heritability, selection, and evolutionary constraints. *Evolution*, **64**, 2614–2628.
- Johnson DW, Christie MR, Moye J, Hixon MA (2011) Genetic correlation between adults and larvae in a marine fish: potential effects of fishery selection on population replenishment. *Evolutionary Applications*, **4**, 621–633.
- Johnson DW, Grorud-Colvert K, Sponaugle S, Semmens BX (2014) Phenotypic variation and selective mortality as major drivers of recruitment variability in fishes. *Ecology Letters*, 17, 743–755.
- Jones GP, Milicich MJ, Emslie MJ, Lunow C (1999) Self-recruitment in a coral reef population. *Nature*, 402, 802–804.
- Jones GP, Planes S, Thorrold SR (2005) Coral reef fish larvae settle close to home. *Current Biology*, **15**, 1314–1318.
- Jost L (2008) G_{ST} and its relative do no measure differentiation. Molecular Ecology, **17**, 4015–4026.
- Knapp RA, Warner RR (1991) Male parental care and female choice in the bicolor damselfish, *Steastes partitus*: bigger is not always better. *Animal Behaviour*, **41**, 747–756.
- Kritzer JP, Sale PF (2004) Metapopulation ecology in the sea: from Levins' model to marine ecology and fisheries science. *Fish and Fisheries*, **5**, 131–140.
- Lacson JM (1992) Minimal genetic variation among samples of six species of coral reef fishes collected at La Parguera, Puerto Rico, and Discovery Bay, Jamaica. *Marine Biology*, **112**, 327–331.
- Leis JM, McCormick MI (2002) The biology, behavior, and ecology of the pelagic, larval stage of coral reef fishes. In: *Coral Reef Fishes. Dynamics and Diversity in a Complex Ecosystem* (ed. Sale PF), pp. 171–199. Academic Press, San Diego, California.
- Lester SE, Ruttenberg BI (2005) The relationship between pelagic larval duration and range size in tropical reef fishes: a synthetic analysis. *Proceedings of the Royal Society B*, **272**, 585– 591.
- Levin LA (2006) Recent progress in understanding larval dispersal: new direction and digressions. *Integrative and Comparative Biology*, 46, 282–297.
- Levitan DR (2005) The distribution of male and female reproductive success in a broadcast spawning marine invertebrate. *Integrative and Comparative Biology*, **45**, 848–855.
- Lipcius RN, Stockhausen WT, Eggleston DB, Marshal LS Jr, Hickey B (1997) Hydrodynamic decoupling of recruitment, habitat quality and adult abundance in the Caribbean spiny lobster: source-sink dynamics? *Marine and Freshwater Research*, **48**, 807–815.
- Lipcius RN, Stockhausen WT, Eggleston DB (2001) Marine reserves for Caribbean spiny lobster: empirical evaluation and theoretical metapopulation recruitment dynamics. *Marine and Freshwater Research*, **52**, 1589–1598.

- McCook LJ, Almany GR, Berumen ML *et al.* (2009) Management under uncertainty: guide-lines for incorporating connectivity into the protection of coral reefs. *Coral Reefs*, **28**, 353–366.
- Meirmans PG, Hedrick PW (2011) Assessing population structure: F_{ST} and related measures. *Molecular Ecology Resources*, 11, 5–18.
- Meirmans PG, Van Tiernderen PH (2004) GENOTYPE and GENO-DIVE: two programs for the analysis of genetic diversity of asexual organisms. *Molecular Ecology Resources*, **4**, 792–794.
- Miller JA, Shanks AL (2004) Evidence for limited larval dispersal in black rockfish (*Sebastes melanops*): implications for population structure and marine-reserve design. *Canadian Journal of Fisheries and Aquatic Sciences*, 61, 1723–1735.
- Myrberg AA Jr (1972) Social dominance and territoriality in the bicolor damselfish, *Eupomacentrus partitus* (Poey). *Behaviour*, **41**, 207–231.
- Ospina-Guerrero SP, Landinez-Garcia RM, Rodriguez-Castro DJ, Arango R, Marquez E (2008) Genetic connectivity of *Stegastes partitus* in the south Caribbean evidenced by microsatellite analysis. *Ciencias Marinas*, **3**, 155–163.
- Palumbi SR (2003) Population genetics, demographic connectivity, and the design of marine reserves. *Ecological Applications*, **13**, S146–S158.
- Planes S, Jones GP, Thorrold SR (2009) Larval dispersal connects fish populations in a network of marine protected areas. Proceedings of the National Academy of Sciences of the United States of America, 106, 5693–5697.
- Purcell JFH, Cowen RK, Hughes CR, Williams DA (2009) Population structure in a common Caribbean coral-reef fish: implications for larval dispersal and early life-history traits. *Journal of Fish Biology*, **74**, 403–417.
- R Core Team (2012) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. Available from http://www.R-project.org. ISBN 3-900051-07-0.
- Raymond M, Rousset F (1995) GENEPOP, version 1.2: population genetics software for exact tests and ecumenicism. *Journal of Heredity*, **86**, 248–249.
- Rousset F (2008) GENEPOP'007: a complete reimplementation of the GENEPOP software for Windows and Linux. *Molecular Ecology Resources*, **8**, 103–106.
- Saenz-Agudelo P, Jones GP, Thorrold SR, Planes S (2012) Patterns and persistence of larval retention and connectivity in a marine fish metapopulation. *Molecular Ecology*, **21**, 4695–4705.
- Salas E, Molina-Urena H, Walter RP, Heath DD (2010) Local and regional genetic connectivity in a Caribbean coral reef fish. *Marine Biology*, **157**, 437–445.
- Sambrook J, Russell DW (2001) *Molecular Cloning: A Laboratory Manual*, 3rd edn. Cold Springs Harbor Laboratory Press, Cold Springs Harbor, New York.
- Schmale MC (1981) Sexual selection and reproductive success in males of the bicolor damselfish, *Eupomacentrus partitus* (Pisces: Pomacentridae). *Animal Behaviour*, **29**, 1172– 1184.
- Selkoe KA, Gaines SD, Caselle JE, Warner RR (2006) Current shifts and kin aggregation explain genetic patchiness in fish recruits. *Ecology*, 87, 3082–3094.
- Shanks AL (2009) Pelagic larval duration and dispersal distance revisited. *Biological Bulletin*, 216, 373–385.
- Siegel DA, Mitarai S, Costello CJ et al. (2008) The stochastic nature of larval connectivity among nearshore marine popu-

3408 T. J. PUSACK ET AL.

lations. Proceedings of the National Academy of Sciences of the United States of America, **105**, 8974–8979.

- Sponaugle S, Cowen RK (1996) Larval supply and patterns of recruitment for two Caribbean reef fishes, *Stegastes partitus*, and *Acanthurus bahianus*. *Marine and Freshwater Research*, **47**, 433–447.
- Stockhausen WT, Lipcius RN (2001) Single large or several small marine reserves for the Caribbean spiny lobster? *Marine and Freshwater Research*, **52**, 1605–1614.
- Stockhausen WT, Lipcius RN, Hickey BM (2000) Joint effects of larval dispersal, population regulation, marine reserve design, and exploitation on production and recruitment in the Caribbean spiny lobster. *Bulletin of Marine Science*, 66, 957–990.
- Swearer SE, Caselle JE, Lea DW, Warner RR (1999) Larval retention and recruitment in an island population of a coral-reef fish. *Nature*, **402**, 799–802.
- Thiessen RJ, Heath DD (2007) Characterization of one trinucleotide and six dinucleotide microsatellite markers in bicolor damselfish, *Stegastes partitus*, a common coral reef fish. *Conservation Genetics*, 8, 983–985.
- Tracey ML, Bellet NF, Gravem CD (1975) Excess allozyme homozygosity and breeding population structure in the mussel *Mytilus californianus*. *Marine Biology*, **32**, 303–311.
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, **4**, 535–538.
- Victor BC, Wellington GM (2000) Endemism and the pelagic larval duration of reef fishes in the eastern Pacific Ocean. *Marine Ecology Progress Series*, **205**, 241–248.
- Villegas-Sanchez CA, Rivera-Madrid R, Arias-Gonzales JE (2010) Small-scale genetic connectivity of bicolor damselfish (*Stegastes partitus*) recruits in Mexican Caribbean reefs. *Molecular Ecology*, **21**, 4672–4680.
- Wahlund S (1928) Zusammensetzun von Populationen und Korrelation-sersheinungen von Standpunkt der Verebungslehre aus betrachtet. *Hereditas*, **11**, 65–106.
- Waples RS (1998) Separating the wheat from the chaff: patterns of genetic differentiation in high gene flow species. *Journal of Heredity*, 89, 438–450.
- Watson JR, Kendall BE, Siegel DA, Mitarai S (2012) Changing seascapes, stochastic connectivity, and marine metapopulation dynamics. *The American Naturalist*, **180**, 99–112.
- Weersing K, Toonen RJ (2009) Population genetics, larval dispersal, and connectivity in marine systems. *Marine Ecology Progress Series*, **393**, 1–12.
- Weir BS, Cockerham CC (1984) Estimating F-Statistics for the analysis of population structure. *Evolution*, 38, 1358–1370.
- Wellington GM, Victor BC (1989) Planktonic larval duration of one hundred species of Pacific and Atlantic damselfishes (Pomacentridae). *Marine Biology*, **101**, 557–567.
- Werner FE, Cowen RK, Paris CB (2007) Coupled biological and physical models. Oceanography, 20, 54–69.

- White C, Selkoe KA, Watson J *et al.* (2010) Ocean currents help explain population genetic structure. *Proceedings. Biological Sciences/The Royal Society*, **277**, 1685–1694.
- Williams DA, Purcell JFH, Hughes CR, Cowen RK (2003) Polymorphic microsatellite loci for population studies of the bicolor damselfish, *Stegastes partitus* (Pomacentridae). *Molecular Ecology Notes*, **3**, 547–549.

C.D.S, D.W.J., M.A.H, and M.R.C planned the study. C.D.S, D.W.J., M.R.C, M.A.H, and T.J.P conducted field work. M.R.C. and T.J.P conducted the laboratory work. T.J.P. analysed the data, and wrote the paper with guidance by M.R.C., with all authors contributing to revisions.

Data accessibility

All microsatellite genotypes and R scripts have been deposited in the Dryad Digital Repository (doi:10.5061/dryad.qr7f5).

Supporting information

Additional supporting information may be found in the online version of this article.

Table S1 Pairwise F_{ST} values for adult vs. adult bicolor damselfish populations (above diagonal) and their corresponding *P*-values (below diagonal).

Table S2 Pairwise G'_{ST} values for both adult vs. adult (above diagonal) and recruit vs. recruit (below diagonal) bicolor damselfish populations.

Table S3 Pairwise G''_{ST} values for both adult vs. adult (above diagonal) and recruit vs. recruit (below diagonal) bicolor damselfish populations.

Table S4 Pairwise D_{est} values for both adult vs. adult (above diagonal) and recruit vs. recruit (below diagonal) bicolor damselfish populations.

Table S5 Pairwise F_{ST} values among all adult vs. recruit populations. Black boxes with white text indicate significant F_{ST} values based on a sequential bonferroni correction.

Table S6 Pairwise F_{ST} values recruit vs. recruit bicolor damselfish populations (below diagonal) and their corresponding *P*-values (above diagonal).